

Phenomenology of the 0.7 conductance feature

D. J. Reilly,^{*} Y. Zhang and L. DiCarlo

Department of Physics, Harvard University, Cambridge, 02138, USA

Abstract

We describe a phenomenological model for the conductance feature near $0.7 \times 2e^2/h$ that occurs in quantum point contacts. We focus on the transconductance at finite source-drain bias and contrast our model with the results expected from a single-particle picture. Good agreement is seen in comparing the model with experimental data, taken on ultra-low-disorder GaAs induced electron systems. Although simple, our phenomenology suggests important boundary conditions for an underlying microscopic theory.

Key words: 0.7, QPC, phenomenological, quantum wire, conductance

PACS:

1. Introduction

The conductance of a quantum point contact (QPC), formed via geometric or electrostatic confinement of a two-dimensional electron system (2DES), is quantized in units of $2e^2/h$ [1,2]. This quintessentially quantum-mechanical phenomena is well explained in terms of the single-particle discrete energy spectrum associated with the confining potential [3]. QPCs also exhibit unexpected non-quantized features that cannot be explained within this non-interacting, single-particle picture. A clear example is the feature occurring below the first conductance plateau, in the range $0.5 - 0.7 \times 2e^2/h$ and at equivalent positions for higher plateaus. This anomalous conductance structure was observed in the earliest transport measurements on ballistic QPCs [1,2] but has remained largely unexplained despite numerous experimental investigations [4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19] and is re-

garded as a key outstanding problem in mesoscopic physics [20].

A prime result, initially uncovered by Thomas *et al.*, [4] is the evolution of the 0.7 *feature* with increasing parallel magnetic field into the well understood Zeeman spin-split plateau at $0.5 \times 2e^2/h$. This observation strongly suggests that the 0.7 structure at $B = 0$ is related to a many-body state in which symmetry is broken between spin-up and spin-down electrons.

2. Phenomenological model

Here we describe a simple phenomenological model that is in striking agreement with the experimental behavior of the conductance feature with temperature T , magnetic field B , source-drain bias V_{sd} and potential profile. At the core of this model is the conjecture that a density-dependent spin-gap opens in the one-dimensional (1D) energy spectrum as the gate voltage is made less negative [13,21]. Fig. 1a captures the

^{*} Email: reilly@fas.harvard.edu




Fig. 1. a) Energy level configuration underlying our phenomenological model. A spin-gap opens in the 1D energy spectrum as a function of gate bias V_g . The Fermi-level E_F is parabolic with gate voltage because of the singularity in the 1D density of states. b) Shows the *difference* between E_F and the spin-up and spin-down band-edges. c) Shows conductance calculations based on our model (red and dashed line) together with experimental data (black and solid line) taken at $T=50\text{mK}$. d) Shows calculated conductance in the context of the model for increasing parallel magnetic field B . Consistent with experiment we observe a smooth evolution from 0.7 to $0.5 \times 2e^2/h$. Panels c) and d) are taken from Reference [21].

essence of the model. Before the first 1D sub-band is occupied (below energy E_1 in Fig. 1a) the spin-up and spin-down sub-bands are degenerate. As the 1D states begin to fill the spin-down band rapidly moves down in energy with the available spin-up states moving higher in energy above the Fermi-level (E_F). The model assumes a parabolic dependence of the Fermi-energy on gate voltage ($E_F \sim V_g^2$) in connection with the usual $E^{-1/2}$ form of the 1D density of states [22]. For a detailed discussion of this model we refer the reader to Refs. [21,16].

Features in conductance are related to the occupation probability of the 1D sub-bands and are a function of temperature and the energy difference between the sub-band edge and E_F . Within the context of our model, Fig. 1b shows the difference between the spin-dependent band-edge energy and E_F as a function of gate voltage V_g . It is the functional form of this *difference* between the spin band-edges and E_F that relates to the experimental data (since it sets the conductance) and of prime importance in suggesting possible ‘boundary conditions’ of an underlying microscopic

theory. We point out that in contrast to a similar model based on pinning of the band-edge [23], in our picture the spin-down energy continues to move rapidly below the Fermi-level as the gate voltage is swept.

Figure 1c (taken from Ref [21]) compares differential conductance $g = di/dv$ calculations based on this model with data taken on an ultra-low-disorder quantum wire at $T=50\text{mK}$ (See Refs [13,16,21] for experimental details). We take the simplest approximation and calculate conductance with a step transmission function in the limit of zero source-drain bias. Importantly, the only free-parameter needed to fit the model to the experimental data is the rate at which the spin-gap opens with gate voltage $\gamma = dE_{\uparrow\downarrow}/dV_g$ [21]. Note the small discrepancy between model and experiment at the top right of Fig. 1c which is due to the step transmission function (no tunneling) that is used in the calculation rather than a more realistic smoothly varying $T(E)$ [3]. Fig. 1d (also taken from Ref. [21]) shows conductance calculations as a function of increasing parallel magnetic field, $B_{\parallel}$ in which a smooth evolution of the conductance feature from 0.7 to $0.5 \times 2e^2/h$ is observed, consistent with the experimental results [4,15,24].

3. Single-particle results

We now focus on the transconductance (dg/dV_g) at finite V_{sd} which facilitates the study of transitions between the conductance plateaus as a function of chemical potential or gate voltage V_g . We firstly compare transconductance plots based on our model to the single-particle case at zero and finite magnetic field. Fig. 2a shows a transconductance intensity plot as a function of V_{sd} and V_g calculated using the usual single-particle formalism [25,26]. We assume the conductance at finite bias is approximated by the weighted average of two zero V_{sd} conductances, one for the chemical potential of the source ($E_F + \beta eV_{sd}$) and the other for the drain ($E_F - (1 - \beta)eV_{sd}$), where β characterizes the symmetry of the potential drop across the constriction [26]. Centered about $V_{sd}=0$ are the linear response integer-plateau diamonds (dark) at $0, 1$ and $2 \times 2e^2/h$. Symmetric either side of the integer plateaus are the finite bias ‘half-plateaus’ that occur when the chemical potential of the source (μ_s)




Fig. 4. Transconductance calculations based on the model for $V_{sd}=0$ as a function of gate bias V_g and in-plane magnetic field, B (traces are off-set). We use a bulk value for the electron g -factor of 0.44. See main text and Ref. [24] for detailed discussion.

of high bias 1.25 features provides further evidence that a spin dependent energy gap remains open well below the Fermi-level, continuing to increase (slightly) as the 1D density grows and higher sub-bands are populated.

5. 0.7 analogs

We now discuss our phenomenological model in the context of a recent experiment by Graham *et al.*, [24], in which the sub-band energies can be made to cross at very high in-plane magnetic fields. In addition to the usual evolution of the 0.7 feature towards $0.5 \times 2e^2/h$, Graham *et al.*, discovered the appearance of similar conductance structure at high $B_{||}$ evolving from $1.5 \times 2e^2/h$ into the plateau at $2e^2/h$. This feature is termed a ‘0.7-analog’. In Fig. 4 we show transconductance calculations based on our model as a function of increasing Zeeman energy (bottom to top). For simplicity we ignore any changes in the electron g -factor related to correlation or diamagnetic effects, taking the bulk value of 0.44. This likely accounts for the larger magnetic fields required in our model to achieve sub-band crossing in comparison to experimental observation.

Consistent with the experimental results of Graham *et al.*, [24] we observe the usual 0.7 feature at $B=0$, indicated in Fig. 4 by the black region separating the spin-up and spin-down transconductance lines that evolve away from each other as B is increased. Without adding any new parameters to our model we also find good

agreement with the experimental results at magnetic fields where the $N=1$ spin-up sub-band crosses the $N=2$ spin-down sub-band. As in the case of the 0.7 structure at $B=0$, our model assumes that at high $B_{||}$ as each spin-band is populated, the Fermi energy E_F is proportional to V_g^2 and the gate-dependent spin gap (intrinsic) adds to the (extrinsic) Zeeman energy of the B -field.

Following the crossing of sub-bands there is a discontinuous shift in δV_g from the crossing point, marking the appearance of the 0.7 analog [24]. Our calculations also account for the appearance of higher order 0.7 analogs at the crossings of $N=2$ spin-up with $N=3$ spin-down sub-bands. The temperature dependence of the 0.7 analogs follows the usual activated behavior of the 0.7 feature in our model (not shown). Very recent numerical work [28] using spin-density functional theory also shows good agreement with the experimental results of Graham *et al.*. How the functional form of our phenomenology relates to these more detailed calculations is an interesting question for future work.

The strong agreement between the transconductance calculations and the experimental data of Graham *et al.*, provide further evidence that this model captures the essence of the functional form underlying the 0.7 feature. Despite this agreement however, the observation of a zero bias anomaly in the bias spectroscopy and the low temperature *restoration* of the conductance as observed by Cronenwett *et al.*, [15] cannot be explained without extending this model to also include Kondo spin screening in the regime $T < T_K$, where T_K is the Kondo temperature. Perhaps a complete explanation will account for a density-dependent gap that is screened by the formation of a Kondo-like state at low temperatures and bias.

6. Conclusions

We have described a simple phenomenological model for the $0.7 \times 2e^2/h$ conductance feature that occurs in quantum point contacts. Our focus has been on the transconductance at finite source-drain bias which emphasizes transitions between the conductance plateaus. The calculated results based on our model agree well with data taken on both ultra-low-disorder *induced* devices and more traditional split-gated heterostruc-

tures. Without the inclusion of additional parameters this model also accounts for the observation of 0.7 analogs at high magnetic field. Although largely empirical, we believe the functional form underlying this simple picture may prove useful in uncovering a detailed microscopic theory of this effect.

7. Acknowledgments

The authors wish to thank C. M. Marcus, T. M. Buehler, D. T. McClure, J. L. O'Brien, S. Das Sarma, K.-F. Berggren, M. J. Biercuk, R. G. Clark, A. Dzurak and A. R. Hamilton for useful discussions. D.J.R. is indebted to L. N. Pfeiffer and K. W. West for the excellent heterostructure material used in the experiments.

References

- [1] B. J. van Wees et al. *Phys. Rev. Lett.*, 60:848, 1988.
- [2] D. A. Wharam et al. *J. Phys. C*, 21:L209, 1988.
- [3] M. Büttiker. *Phys. Rev. B*, 41:7906, 1990.
- [4] K. J. Thomas et al. *Phys. Rev. Lett.*, 77:135, 1996.
- [5] K. J. Thomas et al. *Phys. Rev. B*, 58:4846, 1998.
- [6] B. E. Kane et al. *Appl. Phys. Lett.*, 72:3506, 1998.
- [7] A. Kristensen et al. *Physica B*, 249:180, 1998.
- [8] A. Kristensen et al. *Phys. Rev. B*, 62:10950, 2000.
- [9] S. Nuttinck et al. *Jpn. J. Appl. Phys.*, 39:L655, 2000.
- [10] K. J. Thomas et al. *Phys. Rev. B*, 61:R13365, 2000.
- [11] C.-T. Liang et al. *Phys. Rev. B*, 60, 1999.
- [12] K. S. Pyshkin et al. *Phys. Rev. B*, 62:15842, 2000.
- [13] D. J. Reilly et al. *Phys. Rev. B*, 63:R121311, 2001.
- [14] N. J. Appleyard et al. *Phys. Rev. B*, 62:R16275, 2000.
- [15] S. M. Cronenwett et al. *Phys. Rev. Lett.*, 88:226805, 2002.
- [16] D. J. Reilly et al. *Phys. Rev. Lett.*, 89:246801, 2002.
- [17] Jonathan. P. Bird and Yuichi Ochiai. *Science*, 303:1621, 2004.
- [18] P. Roche et al. *Phys. Rev. Lett.*, 93:116602, 2004.
- [19] R. de Picciotto et al. *Phys. Rev. Lett.*, 92:036805, 2004.
- [20] R. Fitzgerald. *Phys. Today*, 55(5):21, 2002.
- [21] D. J. Reilly. *Phys. Rev. B*, 72:033309, 2005.
- [22] K. Hirose and N. S. Wingreen. *Phys. Rev. B*, 64:073305, 2001.
- [23] A. Kristensen and H. Bruus. *Physica Scripta: Proc. 19th Nordic Semiconductor Meeting, 2001*, 2002.
- [24] A. C. Graham et al. *Phys. Rev. Lett.*, 91:136404, 2003.
- [25] L. I. Glazman and A. V. Khaetskii. *Europhys. Lett.*, 9:263, 1989.
- [26] L. Martín-Moreno et al. *J. Phys. C*, 4:1323, 1992.
- [27] N. K. Patel et al. *Phys. Rev. B*, 44:R10973, 1991.
- [28] Karl-Fredrik Berggren, Peter Jaksch, and Irina Yakimenko. *Phys. Rev. B*, 71:115303, 2005.